



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MISCELLANEOUS.

104. Proposed by HARRY S. VANDIVER, Bala, Pa.

A Theorem of Fermat. The area of a right angled triangle with commensurable sides cannot be a square number. [Cf. Chrystal's *Algebra*, Vol. II., page 535.]

Solution by L. C. WALKER, A. M., Professor of Mathematics, Petaluma High School, Petaluma, Cal.

Since the sides of the right-angled triangle may have a common factor, we let af , bf , and cf represent its sides, where a , b , c are prime to each other. Then we have $a^2 + b^2 = c^2$(1).

For the area, $\frac{1}{2}abf^2$, of the triangle to be a square, $\frac{1}{2}ab$ must be a square.

In (1) either a or b must be even and the other odd, because the sum of two odd squares can not be a square. Assume a even and b odd. Then for $\frac{1}{2}ab$ to be a square, b must be a square, and a must be double a square.

Let $a=2m^2$, $b=n^2$, and substitute these values in (1); then we have

$$(2m^2)^2 + (n^2)^2 = c^2$$
.....(2).

Now set $2m^2=2pq$, and $n^2=p^2-q^2$; or

$$n^2 + q^2 = p^2$$
.....(3).

In (3) p is odd, and we will assume q even. Let $q=2a\beta$, $p=a^2+\beta^2$, and we find $m=2a\beta(a^2+\beta^2)$(4).

For $2a\beta$ to be a square, we will assume $a=2m_1^2$ and $\beta=n_1^2$. Substituting these values in (4), we obtain

$$m^2=4m_1^2n_1^2(4m_1^4+n_1^4)$$
.....(5).

In order that the right member of (5) may be a square, we must have

$$4m_1^4+n_1^4=c_1^2,$$

say; which is of the same form of (2). But c_1^2 is less than c^2 . Proceeding in exactly the same way, we can reduce c^2 indefinitely. By our hypothesis c^2 can not be reduced to zero nor less. Hence, results *Fermat's Theorem*.

PROBLEMS FOR SOLUTION.

ARITHMETIC.

154. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

Suppose there is a meadow of 8 acres in which the grass grows uniformly, and that 21 oxen could eat up the whole pasture in 6 weeks, or 18 oxen in 9 weeks; what number of oxen diminished by the removal of 9, at the end of 14 weeks, could eat it up in 18 weeks?